

# MarinTrust Standard V2

# By-product Fishery Assessment Plaice (*Pleuronectes platessa*), FAO 27, ICES 7.e (Western English Channel)

#### MarinTrust Programme

Unit C, Printworks 22 Amelia Street London SE17 3BZ

E: standards@marin-trust.com

T: +44 2039 780 819



# Table 1 Application details and summary of the assessment outcome

|   | Species:                          | Plaice (Pleuronectes platessa)      |  |  |
|---|-----------------------------------|-------------------------------------|--|--|
| Fishery Under<br>Assessment             | Geographical area:                | FAO 27 – Northeast Atlantic         |  |  |
|   | Country of origin of the product: | Denmark , France, Belgium, UK       |  |  |
|   | Stock:                            | ICES 7. e (Western English Channel) |  |  |
| Date                                    | May 2024                          |                                     |  |  |
| Report Code                             | DNK22                             |                                     |  |  |
| Assessor                                | Blanca Gonzalez                   |                                     |  |  |
| Country of origin of the product - PASS | Denmark, France, Belgium, UK      |                                     |  |  |
| Country of origin of the product - FAIL | None                              |                                     |  |  |

| Application details and summary of the assessment outcome |            |                    |                                      |  |  |
|---|------------|--------------------|--------------------------------------|--|--|
| Company Name(s): FF                                       | Skagen A/S |                    |                                      |  |  |
| Country: Denmark  |            |                    |                                      |  |  |
| Email address:  |            | Applicant Code     | e:                                   |  |  |
| <b>Certification Body Deta</b>                            | ails       |                    |                                      |  |  |
| Name of Certification Body:                               |            | LRQA               |                                      |  |  |
| Assessor Peer Reviewer                                    |            | Assessment<br>Days | Initial/Surveillance/<br>Re-approval |  |  |
| Blanca Gonzalez Sam Peacock                               |            | 0.5                | Re-approval                          |  |  |
| Assessment Period May 2024 – May 2025                     |            |                    |                                      |  |  |

| Scope Details                            |  |
|--|--|
| Main Species                             | Plaice (Pleuronectes platessa)                   |
| Stock                                    | ICES 7. e (Western English Channel)              |
| Fishery Location                         | FAO 27 – Northeast Atlantic                      |
| Management Authority<br>(Country/ State) | EU and UK  |
| Gear Type(s)                             | Beam trawl, otter trawl, fixed nets, other gears |
| Outcome of Assessment                    |  |
| Peer Review Evaluation                   | Agree with assessment outcome                    |
| Recommendation                           | PASS   |



### Table 2. Assessment Determination

#### **Assessment Determination**

Plaice (*Pleuronectes platessa*) was assessed as a category C species considering that it is a Least Concern species by the IUCN, it is not in included in any CITES Appendixes, and the stock is managed using biennial quotas relative to established reference points.

Plaice in ICES Division 7. e (western English Channel) is subject to a biennial stock assessment by ICES working group for the Celtic Seas Ecoregion (WGCSE). The last assessment was published in June 2022 based on the rfb rule which uses catches data. Fishing pressure on the stock is above F<sub>MSY</sub> proxy, and the stock-size index is above MSY B<sub>trigger</sub> (I<sub>trigger</sub>). Therefore, both clauses in the assessment were met.

The plaice by-product meets the Marin Trust requirements and it should be re-approved for use as a raw material.

#### **Fishery Assessment Peer Review Comments**

The peer reviewer agrees that this species is eligible for assessment under the MarinTrust byproduct assessment methodology, and that the stock falls into Category C. The most recent stock assessment was adequate to meet the requirements of C1.1, and biomass is currently estimated to be above the target reference point level, meeting the requirements of C1.2. Overall, the peer reviewer agrees that this stock should be approved as a source of byproduct raw material for MarinTrust certified facilities.

| Notes for On-site Auditor |  |  |
|---------------------------|--|--|
| None                      |  |  |
|                           |  |  |
|                           |  |  |
|                           |  |  |
|                           |  |  |
|                           |  |  |
|                           |  |  |
|                           |  |  |
|                           |  |  |
|                           |  |  |



# **Species Categorisation**

**NB:** If any species is categorised as Endangered or Critically Endangered on the IUCN Red List, or if it appears in CITES Appendix 1, it **cannot** be approved for use as an MarinTrust raw material.

#### **IUCN Red list Category**

By-product material from a species listed by IUCN (the International Union for Conservation of Nature) under the Red List for the following categories shall immediately fail the assessment;

- EXTINCT (E) AND EXTINCT IN THE WILD (EW)
- CRITICALLY ENDANGERED (CR) facing an extremely high risk of extinction in the wild.
- ENDANGERED (EN) facing a very high risk of extinction in the wild.

By-product material may be used from the following categories provided that all clauses in the MarinTrust standard are passed.

- VULNERABLE (VU) facing a high risk of extinction in the wild.
- NEAR THREATENED (NT) does not qualify for above now, but is close or is likely to qualify for, a threatened category in the near future.
- LEAST CONCERN (LC) Widespread and abundant.
- DATA DEFICIENT (DD) and NOT EVALUATED (NE)

# **Table 3 Species Categorisation Table**

| Common name | Latin name               | Stock  | Management | Category | IUCN Red List<br>Category <sup>1</sup> | CITES Appendix 1 <sup>2</sup> |
|-------------|--------------------------|--|------------|----------|--|-------------------------------|
| Plaice      | Pleuronectes<br>platessa | ICES 7. e<br>(Western<br>English<br>Channel) | Yes        | С        | Least Concern <sup>3</sup>             | No                            |

<sup>&</sup>lt;sup>1</sup> https://www.iucnredlist.org/

<sup>&</sup>lt;sup>2</sup> https://cites.org/eng/app/appendices.php

<sup>3</sup> https://www.iucnredlist.org/species/135690/50018800



#### **CATEGORY C SPECIES**

In a by-product assessment, Category C species are those which are subject to a species-specific management regime and are usually targeted species in fisheries for human consumption.

Clause C1 should be completed for each Category C species. If there are no Category C species in the fishery under assessment, this section can be deleted. Where a species fails this Clause, it should be assessed as a Category D species instead.

| Spe       | cies  | Name         | Plaice (Pleuronectes platessa)  |      |  |  |
|-----------|---|--------------|---|------|--|--|
| <b>C1</b> | Category C Stock Status - Minimum Requirements  |              |   |      |  |  |
| CI        | C1.1  | Fishery remo | ovals of the species in the fishery under assessment are included in the stock assessment | PASS |  |  |
|           | process, OR are considered by scientific authorities to be negligible.                                      |              |   |      |  |  |
|           | C1.2 The species is considered, in its most recent stock assessment, to have a biomass above the limit PASS |              |   |      |  |  |
|           |   | reference po | int (or proxy), OR removals by the fishery under assessment are considered by scientific  |      |  |  |
|           | authorities to be negligible.   |              |   |      |  |  |
|           |   |              | Clause outcome:   | PASS |  |  |

C1.1 Fishery removals of the species in the fishery under assessment are included in the stock assessment process, OR are considered by scientific authorities to be negligible.

The clause is met considering that:

The plaice in the western English Channel (ICES Division7.e) most recent assessment was published in June 2022 by The International Council for exploration of the Sea (ICES) Working Group for the Celtic Seas Ecoregion (WGCSE). The 2022 assessment is still in force since ICES gave biennial advice for 2023 and 2024; therefore, no new assessment / advice was produced at WGCSE 2023 (ICES 2023). The assessment for this stock is basses on the rfb rule, since this stock is a category 3 data-limited stock, which requires commercial catch (international landings and discards, length frequencies from catch sampling) and one survey index (ICES 2022). Thus, removals of the species are included in the stock assessment process (figure 1).




Figure 1. Plaice catches in Division 7.e (Western English Channel) since 1980 (ICES 2022).



C1.2 The species is considered, in its most recent stock assessment, to have a biomass above the limit reference point (or proxy), OR removals by the fishery under assessment are considered by scientific authorities to be negligible.

The Clause is met considering that:

The 2022 plaice assessment indicates that fishing pressure on the stock is above F<sub>MSY proxy</sub> (figure 1), and the stock-size index is above MSY B<sub>trigger</sub> (I<sub>trigger</sub>) (ICES 2022) (figure 2). ICES advises that when the MSY approach is applied, catches should be no more than 1,219 tonnes in each of the years 2023 and 2024 (ICES 2023).




Figure 1. Plaice in Division 7.e (Western English Channel). Length indicator (mean length of fish in the catch and MSY proxy reference length L  $_{F=M}$ ). The exploitation status is below  $F_{MSY\ proxy}$  when the indicator ratio value is higher than 1 (shown by a blue line) (ICES 2022).




Figure 2. Stock biomass index for plaice in Division 7.e (Western English Channel) above MSY  $B_{trigger}$  ( $I_{trigger}$ ). The horizontal orange lines indicate the average of the biomass index for 2017 to 2019 and for 2020 to 2021 (ICES 2022).

#### References

ICES (2023). Working Group for the Celtic Seas Ecoregion (WGCSE). ICES Scientific Reports. Report. https://doi.org/10.17895/ices.pub.22268980.v1

ICES (2022). Plaice (Pleuronectes platessa) in Division 7.e (western English Channel). ICES Advice: Recurrent Advice. Report. https://doi.org/10.17895/ices.advice.19453631.v1

| Links                      |               |
|----------------------------|---------------|
| MarinTrust Standard clause | 1.3.2.2       |
| FAO CCRF                   | 7.5.3         |
| GSSI                       | D.3.04, D5.01 |



## **CATEGORY D SPECIES**

Category D species are those which are not subject to a species-specific management regime. In the case of mixed trawl fisheries, Category D species may make up the majority of landings. The comparative lack of scientific information on the status of the population of the species means that a risk-assessment style approach must be taken.

| D1     | Species Name   | NA  |                |
|--------|--|---|----------------|
|        | Productivity Attribut  | te Value  | Score          |
|        | Average age at maturity (years)  |   |                |
|        | Average maximum age (years)  |   |                |
|        | Fecundity (eggs/spawning)  |   |                |
|        | Average maximum size (cm)  |   |                |
|        | Average size at maturity (cm)  |   |                |
|        | Reproductive strategy  |   |                |
|        | Mean trophic level   |   |                |
|        |  | Average Productivity Score  |                |
|        | Susceptibility Attribu   | te Value  | Score          |
|        | Availability (area overlap)  |   |                |
|        | Encounterability (the position of the s  | stock/species   |                |
|        | within the water column relative to the  | ne fishing gear)  |                |
|        | Selectivity of gear type   |   |                |
|        | Post-capture mortality   |   |                |
|        |  | Average Susceptibility Score  |                |
|        |  | PSA Risk Rating (From Table D3)   |                |
|        |  | Compliance rating   |                |
|        | Further justification for susceptibility For susceptibility attributes, please pri uncertainty affecting your decision | y scoring (where relevant) ovide a brief rationale for scoring of parameters wher | e there may be |
| Refere | nces   |   |                |
|        |  |   |                |
| Standa | ırd clauses 1.3.2.2  |   |                |



# Table D2 - Productivity / Susceptibility attributes and scores.

| Productivity High productivity attributes (Low risk, score = 1) |                       | Medium productivity<br>(medium risk, score = 2) | Low productivity<br>(high risk, score = 3) |  |
|---|-----------------------|---|--|--|
| Average age<br>at maturity                                      | <5 years              | 5-15 years                                      | >15 years                                  |  |
| Average<br>maximum age  | <10 years             | 10-25 years                                     | >25 years                                  |  |
| Fecundity   | >20,000 eggs per year | 100-20,000 eggs per<br>year                     | <100 eggs per year                         |  |
| Average<br>maximum size   | <100 cm               | 100-300 cm                                      | >300 cm                                    |  |
| Average size<br>at maturity                                     | <40 cm                | 40-200 cm                                       | >200 cm                                    |  |
| Reproductive<br>strategy  | Broadcast spawner     | Demersal egg layer                              | Live bearer                                |  |
| Mean Trophic Level  | <2.75                 | 2.75-3.25                                       | >3.25                                      |  |

| Susceptibility attributes  |   | ow susceptibility<br>ow risk, score = 1)                          | Medium susceptibility<br>(medium risk, score = 2) |   |  | High susceptibility (high risk, score = 3)                             |  |
|--|---|---|---|---|--|--|--|
| Areal overlap<br>(availability)<br>Overlap of the fishing<br>effort with the species<br>range  | <10% overlap  |   | 10-30% overlap                                    |   |  | >30% overlap   |  |
| Encounterability The position of the stock/species within the water column relative to the fishing gear, and the position of the stock/species within the habitat relative to the position of the gear | Low overlap with fishing gear (low encounterability). |   | Medium overlap with fishing gear.                 |   | High overlap with<br>fishing gear (high<br>encounterability).<br>Default score for<br>target species |  |  |
| Selectivity of gear type   | а   | Individuals < size<br>at maturity are<br>rarely caught            | а   | Individuals < size<br>at maturity are<br>regularly caught.                    | а  | Individuals < size<br>at maturity are<br>frequently caught             |  |
| Potential of the gear to<br>retain species   | b   | Individuals < size<br>at maturity can<br>escape or avoid<br>gear. | b   | Individuals < half<br>the size at<br>maturity can<br>escape or avoid<br>gear. | b  | Individuals < half<br>the size at maturity<br>are retained by<br>gear. |  |
| Post-capture mortality (PCM) The chance that, if captured, a species would be released and that it would be in a condition permitting subsequent survival  | re  | vidence of majority<br>leased post-capture<br>Id survival.        | rel   | ridence of some<br>eased post-capture<br>d survival.                          | m  | etained species or<br>ajority dead when<br>leased.                     |  |



| D3                   |             | Average Susceptibility Score |             |          |  |  |
|----------------------|-------------|------------------------------|-------------|----------|--|--|
|                      |             | 1 - 1.75                     | 1.76 - 2.24 | 2.25 - 3 |  |  |
| Average Productivity | 1 - 1.75    | PASS                         | PASS        | PASS     |  |  |
| Score                | 1.76 - 2.24 | PASS                         | PASS        | TABLE D4 |  |  |
|                      | 2.25 - 3    | PASS                         | TABLE D4    | TABLE D4 |  |  |

| <b>D4</b>  | Spe                                  | cies Name  |   |  |  |  |  |  |  |
|--|--------------------------------------|--|---|--|--|--|--|--|--|
| Impacts On Species Categorised as Vulnerable by D1-D3 - Minimum Requirements |                                      |  |   |  |  |  |  |  |  |
|  | D4.1                                 | The potential impacts                                | of the fishery on this species are considered during the management                                 |  |  |  |  |  |  |
|  |                                      | process, and reasonab                                | le measures are taken to minimise these impacts.  |  |  |  |  |  |  |
|  | D4.2                                 | There is no substantia species.                      | There is no substantial evidence that the fishery has a significant negative impact on the species. |  |  |  |  |  |  |
| Outco  | me:                                  |  |   |  |  |  |  |  |  |
| Eviden   |                                      |  |   |  |  |  |  |  |  |
|  | -                                    | ential impacts of the fi<br>easures are taken to mir | shery on this species are considered during the management process, and himise these impacts.       |  |  |  |  |  |  |
| D4.2 T   | here is n                            | o substantial evidence                               | that the fishery has a significant negative impact on the species.                                  |  |  |  |  |  |  |
| Refere   | ences                                |  |   |  |  |  |  |  |  |
| Links  |                                      |  |   |  |  |  |  |  |  |
| Marin <sup>*</sup>   | Frust Standard clause 1.3.2.2, 4.1.4 |  |   |  |  |  |  |  |  |
| FAOC   |                                      |  |   |  |  |  |  |  |  |
| FAO C  | CRF                                  |  | 7.5.1   |  |  |  |  |  |  |