



MarinTrust Standard V2

By-product Fishery Assessment Beaked Redfish (*Sebastes mentella*), FAO 27, ICES 1,2 (Barents Sea Subarea, Norwegian Sea, Spitzbergen, and Bear Island Subarea)

MarinTrust Programme

Unit C, Printworks

22 Amelia Street

London

SE17 3BZ

E: standards@marin-trust.com

T: +44 2039 780 819

Table 1 Application details and summary of the assessment outcome

Fishery Under Assessment	Species:	Beaked redfish (<i>Sebastes mentella</i>)
	Geographical area:	FAO 27 – Northeast Atlantic
	Country of origin of the product:	Norway
	Stock:	ICES 1,2 (Barents Sea Subarea, Norwegian Sea, Spitzbergen, and Bear Island Subarea)
Date	July 2024	
Report Code	FRA64	
Assessor	Blanca Gonzalez	
Country of origin of the product - PASS	Norway	
Country of origin of the product - FAIL	None	

Application details and summary of the assessment outcome			
Company Name(s): Copalis Industrie			
Country: France			
Email address:		Applicant Code:	
Certification Body Details			
Name of Certification Body:		LRQA	
Assessor	Peer Reviewer	Assessment Days	Initial/Surveillance/ Re-approval
Blanca Gonzalez	Sam Peacock	0.5	Surveillance 1
Assessment Period	July 2024 – July 2025		

Scope Details	
Main Species	Beaked redfish (<i>Sebastes mentella</i>)
Stock	ICES 1,2 (Barents Sea Subarea, Norwegian Sea, Spitzbergen, and Bear Island Subarea)
Fishery Location	FAO 27 – Northeast Atlantic
Management Authority (Country/ State)	Joint Norwegian-Russian Fisheries Commission
Gear Type(s)	Pelagic trawl, demersal trawls
Outcome of Assessment	
Peer Review Evaluation	Agree with recommendation
Recommendation	PASS

Table 2. Assessment Determination

Assessment Determination
<p>Beaked redfish (<i>Sebastes mentella</i>) was assessed as a category D species considering that it is a Least Concern species by the IUCN, it is not included in any CITES Appendixes, and since Russian participation in ICES was suspended on March 2022 it has not been possible to run ICES stock assessments or provide ICES advice for the Barents Sea stock as management and data collection for these stocks are shared between Norway and Russia. (ICES 2023).</p> <p>In the Productivity-Susceptibility Analysis (PSA) turbot was awarded an average productivity score of 1.85 and an average susceptibility score of 2, and it passed against Table D3, indicating that beaked redfish is not vulnerable to this fishery.</p> <p>The beaked redfish by-product meets the Marin Trust requirements and it should be remained approved for use as a raw material.</p> <p>ICES (2023). Arctic Fisheries Working Group (AFWG). ICES Scientific Reports. Report. https://doi.org/10.17895/ices.pub.20012675.v1</p>
Fishery Assessment Peer Review Comments
<p>The peer reviewer agrees that this beaked redfish stock should be assessed under Category D. The assessor has completed the PSA correctly and the peer reviewer agrees that the assigned scores result in a PASS outcome on Table D3. For this reason, the byproduct should remain approved for use as a raw material.</p>
Notes for On-site Auditor
<p>There are no concerns that requires attention from the on-site assessor</p>

Species Categorisation

NB: If any species is categorised as Endangered or Critically Endangered on the IUCN Red List, or if it appears in CITES Appendix 1, it **cannot** be approved for use as a MarinTrust raw material.

IUCN Red list Category

By-product material from a species listed by IUCN (the International Union for Conservation of Nature) under the Red List for the following categories shall immediately fail the assessment;

- EXTINCT (E) AND EXTINCT IN THE WILD (EW)
- CRITICALLY ENDANGERED (CR) facing an extremely high risk of extinction in the wild.
- ENDANGERED (EN) facing a very high risk of extinction in the wild.

By-product material may be used from the following categories provided that all clauses in the MarinTrust standard are passed.

- VULNERABLE (VU) facing a high risk of extinction in the wild.
- NEAR THREATENED (NT) does not qualify for above now, but is close or is likely to qualify for, a threatened category in the near future.
- LEAST CONCERN (LC) Widespread and abundant.
- DATA DEFICIENT (DD) and NOT EVALUATED (NE)

Table 3 Species Categorisation Table

Common name	Latin name	Stock	Management	Category	IUCN Red List Category ¹	CITES Appendix 1 ²
Beaked readfish	<i>Sebastes mentella</i>	ICES 1,2 (Barents Sea Subarea, Norwegian Sea, Spitzbergen, and Bear Island Subarea)	No	D	Least Concern ³	No

¹ <https://www.iucnredlist.org/>

² <https://cites.org/eng/app/appendices.php>

³ <https://www.iucnredlist.org/species/154816/115238709>

CATEGORY C SPECIES


In a by-product assessment, Category C species are those which are subject to a species-specific management regime and are usually targeted species in fisheries for human consumption.

Clause C1 should be completed for each Category C species. If there are no Category C species in the fishery under assessment, this section can be deleted. Where a species fails this Clause, it should be assessed as a Category D species instead.

Species Name		NA	
C1	Category C Stock Status - Minimum Requirements		
	C1.1	Fishery removals of the species in the fishery under assessment are included in the stock assessment process, OR are considered by scientific authorities to be negligible.	
	C1.2	The species is considered, in its most recent stock assessment, to have a biomass above the limit reference point (or proxy), OR removals by the fishery under assessment are considered by scientific authorities to be negligible.	
			Clause outcome:
<p>C1.1 Fishery removals of the species in the fishery under assessment are included in the stock assessment process, OR are considered by scientific authorities to be negligible.</p> <p>C1.2 The species is considered, in its most recent stock assessment, to have a biomass above the limit reference point (or proxy), OR removals by the fishery under assessment are considered by scientific authorities to be negligible.</p>			
References			
Links			
MarinTrust Standard clause		1.3.2.2	
FAO CCRF		7.5.3	
GSSI		D.3.04, D5.01	

CATEGORY D SPECIES

Category D species are those which are not subject to a species-specific management regime. In the case of mixed trawl fisheries, Category D species may make up the majority of landings. The comparative lack of scientific information on the status of the population of the species means that a risk-assessment style approach must be taken.

D1	Species Name		Beaked redfish (<i>Sebastes mentella</i>)		
	Productivity Attribute		Value	Score	
	Average age at maturity (years)		7.5 ¹	2	
	Average maximum age (years)		31.6 ¹	3	
	Fecundity (eggs/spawning)		10,247 ¹	2	
	Average maximum size (cm)		77.5 ¹	1	
	Average size at maturity (cm)		25.1 ¹	1	
	Reproductive strategy		Broadcast spawner ¹	1	
	Mean trophic level		4.2 ¹	3	
	Average Productivity Score			1.85	
	Susceptibility Attribute		Value	Score	
	Availability (area overlap)		<10% overlap	1	
	Encounterability (the position of the stock/species within the water column relative to the fishing gear)		High overlap	3	
	Selectivity of gear type		Individuals < size at maturity are rarely caught	1	
	Post-capture mortality		Retained	3	
	Average Susceptibility Score			2	
	PSA Risk Rating (From Table D3)			Pass	
	Compliance rating			Pass	
	Further justification for susceptibility scoring (where relevant)				
	<i>For susceptibility attributes, please provide a brief rationale for scoring of parameters where there may be uncertainty affecting your decision</i>				
	<p>Availability: Beaked redfish distributes in the Eastern Atlantic: Norwegian Sea from Lofoten Island northward to the western and northern coasts of Spitsbergen; southern part of the Barents Sea rarely to 35°E, on the Iceland-Faroes Ridge, Iceland and Greenland. Western Atlantic: Baffin Bay to Nova Scotia in Canada¹, and the ICES 1, 2 regions (Barents Sea Subarea, Norwegian Sea, Spitzbergen, and Bear Island Subarea) only overlaps with less than 10% of the species distribution (figure 1).</p>				
					
	<p>Figure 1. Beaked redfish distribution¹ and in red square the ICES 1, 2 regions under assessment².</p>				

	<p>Encounterability: from 2014 onwards most of the catch is taken as targeted catch³.</p> <p>Selectivity of gear type: catch length size range is from 30 - 42 cm ⁴</p> <p>Post-capture mortality: retained ³.</p>
<p>References</p> <p>1 https://fishbase.se/summary/Sebastes-mentella.html</p> <p>2 https://fish-commercial-names.ec.europa.eu/fish-names/fishing-areas/fao-area-27_en</p> <p>3 ICES (2023). Arctic Fisheries Working Group (AFWG). ICES Scientific Reports. Report. https://doi.org/10.17895/ices.pub.20012675.v1</p> <p>4 ICES (2020). Arctic Fisheries Working Group (AFWG). ICES Scientific Reports. Report. https://doi.org/10.17895/ices.pub.6050</p>	
<p><i>Standard clauses 1.3.2.2</i></p>	

Table D2 - Productivity / Susceptibility attributes and scores.

Productivity attributes	High productivity (Low risk, score = 1)	Medium productivity (medium risk, score = 2)	Low productivity (high risk, score = 3)
Average age at maturity	<5 years	5-15 years	>15 years
Average maximum age	<10 years	10-25 years	>25 years
Fecundity	>20,000 eggs per year	100-20,000 eggs per year	<100 eggs per year
Average maximum size	<100 cm	100-300 cm	>300 cm
Average size at maturity	<40 cm	40-200 cm	>200 cm
Reproductive strategy	Broadcast spawner	Demersal egg layer	Live bearer
Mean Trophic Level	<2.75	2.75-3.25	>3.25

Susceptibility attributes	Low susceptibility (Low risk, score = 1)	Medium susceptibility (medium risk, score = 2)	High susceptibility (high risk, score = 3)
Areal overlap (availability) Overlap of the fishing effort with the species range	<10% overlap	10-30% overlap	>30% overlap
Encounterability The position of the stock/species within the water column relative to the fishing gear, and the position of the stock/species within the habitat relative to the position of the gear	Low overlap with fishing gear (low encounterability).	Medium overlap with fishing gear.	High overlap with fishing gear (high encounterability). Default score for target species
Selectivity of gear type Potential of the gear to retain species	a Individuals < size at maturity are rarely caught	a Individuals < size at maturity are regularly caught.	a Individuals < size at maturity are frequently caught
	b Individuals < size at maturity can escape or avoid gear.	b Individuals < half the size at maturity can escape or avoid gear.	b Individuals < half the size at maturity are retained by gear.
Post-capture mortality (PCM) The chance that, if captured, a species would be released and that it would be in a condition permitting subsequent survival	Evidence of majority released post-capture and survival.	Evidence of some released post-capture and survival.	Retained species or majority dead when released.

D3		Average Susceptibility Score		
		1 - 1.75	1.76 - 2.24	2.25 - 3
Average Productivity Score	1 - 1.75	PASS	PASS	PASS
	1.76 - 2.24	PASS	PASS	TABLE D4
	2.25 - 3	PASS	TABLE D4	TABLE D4

D4 Species Name			
Impacts On Species Categorised as Vulnerable by D1-D3 - Minimum Requirements			
D4.1	The potential impacts of the fishery on this species are considered during the management process, and reasonable measures are taken to minimise these impacts.		
D4.2	There is no substantial evidence that the fishery has a significant negative impact on the species.		
Outcome:			
Evidence D4.1: The potential impacts of the fishery on this species are considered during the management process, and reasonable measures are taken to minimise these impacts. D4.2 There is no substantial evidence that the fishery has a significant negative impact on the species.			
References			
Links			
MarinTrust Standard clause		1.3.2.2, 4.1.4	
FAO CCRF		7.5.1	
GSSI		D.5.01	